

# NON-OPIOID CANCER PAIN THERAPY

Prim dr.sc.med Marijana Milović-Kovačević

Oktober 2019



# CANCER PAIN

Despite guidelines, different types of pain or pain syndromes were present in all stages of cancer and were not adequately treated in a significant percentage of patients, ranging from 56% to 82.3%

*Fallon M et al. Management of cancer pain in adult patients . ESMO Clinical practice guidelines. Ann Oncol 2018*

## CLINICAL PRACTICE GUIDELINES

# Management of cancer pain in adult patients: ESMO Clinical Practice Guidelines<sup>†</sup>

M. Fallon<sup>1</sup>, R. Giusti<sup>2</sup>, F. Aielli<sup>3</sup>, P. Hoskin<sup>4</sup>, R. Rolke<sup>5</sup>, M. Sharma<sup>6</sup> & C. I. Ripamonti<sup>7</sup>, on behalf of the ESMO  
Guidelines Committee\*



26 pages


# ASSESSMENT OF PAIN TO IMPROVE CHOICE OF THERAPY

**Nocioceptive** (ongoing tissue damage)

- Somatic (e.g. bone pain)
- Visceral (e.g. gut or hepatic pain)

**Neuropathic**

- Damage or dysfunction in the nervous system (e.g. brachial plexopathy or spinal cord compression)



# PRACTICAL APPROACH: PAIN ASSESSMENT TOOL


1. What has been your worst pain in the last 24h on a scale 0-10 ?
2. Monitor if the pain is  $< 3$
3. More detailed assessment if the pain is  $> 3$
4. Appropriate analgesic and monitor analgesic side effects



Verbal rating scale	
No pain	1
Very mild	2
Mild	3
Moderate	4
Severe	5
Very severe	6



# ESMO Guideline: Treatment of cancer pain



<sup>a</sup>Do not switch between weak opioids.

BTcP, breakthrough cancer pain; NRS, numerical rating scale; NSAID, nonsteroidal anti-inflammatory drug; t.d., transdermal.

# RECOMMENDATIONS

- Patients should be informed about pain and pain management and should be encouraged to take an active role in their pain management [II, B].
- The onset of pain should be prevented by means of around-the clock (ATC) administration, taking into account the half-life, bioavailability and duration of action of different drugs [II, B].
- Analgesics for chronic pain should be prescribed on a regular basis and not on an 'as required' schedule [V, D].
- The oral route of administration of analgesic drugs should be advocated as the first choice [IV, C].



# PHARMACOLOGIC MANAGEMENT



# TREATING PAIN

## PHARMACOLOGICAL TREATMENTS

- Analgesic (pain relieving) drugs are mainstay of pain control
- Include **central** acting -opioid drugs and **peripherally** acting nonopioid

# NONOPIOIDS

- Generally the first class of drugs used for treatment of pain
- Useful for **acute and chronic** pain from a variety of causes such as: surgery, trauma, arthritis, and cancer
- Have a **ceiling effect** to analgesia which indicates that there is a dose beyond which there is no improvement in the analgesic effect and there may be an increase in side effects
- Does not produce **tolerance or physical** dependence
- Works primarily **at the site of injury**, or peripherally

# MILD PAIN

- Paracetamol (acetaminophen) or a nonsteroidal drug (NSAID) such as ibuprofen or aspirin.
- NSAID have **anti-inflammatory, analgesic and antipyretic** effects.
- The anti-inflammatory action relieves pain by interfering with cyclooxygenase.
- Acetaminofen may be administered as a **single medication or in combination** with other analgesics
- The use of selective NSAIDs designated as selective COX-2 inhibitors have significant cardiovascular and cerebrovascular risks which have limited their utilization

# MILD TO MODERATE PAIN

- Paracetamol, an NSAID and /or paracetamol in a combination product with a **weak opioid** such as hydrocodone, may provide greater relief than their separate use
- NSAIDs block synthesis of ***prostaglandin***
- Examples are salicylates/aspirin/; NSAIDs /ibuprofen/; COX2 inhibitors /celecoxib/;acetaminophen

# ANALGESIC ADJUVANTS

- Are classes of medications that may potentiate the effects of opioids or nonopioids are especially important when treating pain that does not respond well to traditional analgesics alone
- **STEROIDS**-May reduce pain by decreasing inflammation and the resultant compression of healthy tissues
- **BENZODIAZEPINES** –These drugs do not provide pain relief except in the treatment of muscle spasms
- **TRICYCLIC ANTIDEPRESSANTS**-Amitripyline have been shown to relieve pain realated to neuropathy and other painful nerve related conditions-must be taken for days before they are fully effective

# ROUTES FOR ANALGESIC ADMINISTRATION

## ORAL

- Preferred route in most cases, convenient, inexpensive
- Slower onset than IV, can provide consistent blood levels

## RECTAL

- May be used to provide local or systemic pain relief
- Can be used when patient is unable to take oral medication

## TRANSDERMAL PATCH

- For chronic pain, easy to apply, delivers pain relief for 3 days without patch change
- 12-hour delay before effective drug level reached and delay in excreting once removed

## INTRAVENOUS

- Preferred route for postoperative and chronic cancer pain for pt who can not tolerate oral route, provides rapid relief –continuous infusion provides steady drug level; Difficult to use in home care setting

## INTRAMUSCULAR

- For acute pain, rapid pain relief , painful, use only if other routes cannot be used



**PAIN**